

BIOFORS ND-IB 0.25

Generic Name:

BIOFORS ND-IB Inactivated Vaccine

General Description:

BIOFORS ND-IB 0.25 contains inactivated Newcastle Disease Virus (NDV) Ishii Strain, and inactivated Infectious Bronchitis Virus (IBV) M41 Strain, inactivated IBV TS I Strain and inactivated IBV TS II Strain in a stable oil emulsion. NDV Ishii Strain, IBV M41 Strain, IBV TS I Strain and IBV TS II were amplified in embryonated chicken eggs. Allantoic fluids of embryonated chicken eggs containing amplified NDV Ishii Strain, IBV M41 Strain, IBV TS I Strain and IBV TS II Strain were harvested and inactivated. Oil adjuvant was then added. This vaccine has a physical appearance as white oil emulsion.

Composition per Dose (0.25mL):

Newcastle Disease Virus Ishii strain	$\geq 10^{7.7} \text{EID}_{50}$
Infectious Bronchitis Virus M41 strain	$\geq 10^{6.2} \text{EID}_{50}$
Infectious Bronchitis Virus TS I strain	$\geq 10^{6.2} \text{EID}_{50}$
Infectious Bronchitis Virus TS II strain	$\geq 10^{6.2} \text{EID}_{50}$

Indication for Use:

Prevention of Newcastle Disease and Infectious Bronchitis in poultry.

Method of Vaccination:

Administer 1 dose (0.25 mL) to each chicken of 2 weeks old or younger via subcutaneous injection. Administer 1 dose (0.25 mL) to each chicken of 2 weeks old or older via intramuscular injection.

Storage Conditions:

Store in a dark place at 2 to 8 °C. Do not freeze.

Packaging:

200Doses/50mL/Bottle, 400Doses/100mL/Bottle, 500Doses/125mL/Bottle, 1000Doses/250mL/Bottle, and 2000Doses/500mL/Bottle

Shelf Life:

2 years

Precautions:

- Follow the instructions from and use under supervision by a veterinarian.
- Recover vaccine to room temperature before use.
- Shake well to mix thoroughly before use
- Protect from sunlight and high temperatures.
- Vaccinate healthy chicken only.