

BIOFORS ND-IC ISHII 0.25

Generic Name:

BIOFORS ND-IC 0.25 Inactivated Vaccine

General Description:

BIOFORS ND-IC ISHII 0.25 contains inactivated Newcastle Disease Virus (NDV) Ishii Strain and inactivated *Avibacterium paragallinarum* Serotype A (IC-A), inactivated *Avibacterium paragallinarum* Serotype C (IC-C) in a stable oil emulsion. NDV Ishii Strain was amplified in embryonated chicken eggs, whereas IC-A and IC-C were amplified in culture media. Allantoic fluid of embryonated chicken eggs containing amplified NDV Ishii Strain and culture media containing amplified IC-A and IC-C were harvested and inactivated. Oil adjuvant was then added. This vaccine has a physical appearance as white oil emulsion.

Composition per Dose (0.25mL):

Newcastle Disease Virus Ishii strain..... $\geq 10^{8.5}$ EID₅₀
Avibacterium paragallinarum IC-A serotype..... $\geq 10^{7.6}$ CFU
Avibacterium paragallinarum IC-C serotype..... $\geq 10^{7.6}$ CFU

Indication for Use:

Prevention of Newcastle Disease and Infectious Coryza in poultry.

Method of Vaccination:

Administer 1 dose (0.25 mL) to each chicken via intramuscular injection.

Withdrawal Period:

8 weeks

Storage Conditions:

Store in a dark place at 2-8°C. Do not freeze.

Packaging:

200Doses/50mL/Bottle, 400Doses/100mL/Bottle, 1000Doses/250mL/Bottle, and 2000Doses/500mL/Bottle

Shelf Life:

2 years

Precautions:

- Follow the instructions and use under supervision by a veterinarian.
- Recover vaccine to room temperature before use.
- Shake well to mix thoroughly before use.
- Protect from sunlight and high temperatures.
- Vaccinate healthy chickens only.