

BIOFORS ND-IC GEL ISHII 0.5

Generic Name:

BIOFORS ND-IC Inactivated Vaccine

General Description:

BIOFORS ND-IC GEL ISHII 0.5 contains inactivated Newcastle Disease Virus (NDV) Ishii Strain and inactivated Avibacterium paragallinarum Serotype A (IC-A) and inactivated Avibacterium paragallinarum Serotype C (IC-C). NDV Ishii was amplified in embryonated chicken eggs, whereas IC-A and IC-C were amplified in culture media. Allantoic fluids of embryonated chicken eggs containing amplified NDV Ishii and culture media containing amplified IC-A and IC-C were harvested and inactivated. Aluminum hydroxide gel as then added. This vaccine has a physical appearance as yellowish white suspension.

Composition per Dose (0.5mL):

Newcastle Disease Virus Ishii strain	$\geq 10^{8.5} \text{EID}_{50}$
<i>Avibacterium paragallinarum</i> serotype A	$\geq 10^{8.7} \text{CFU}$
<i>Avibacterium paragallinarum</i> serotype C	$\geq 10^{8.7} \text{CFU}$

Indication for Use:

Prevention of Newcastle Disease and Infectious Coryza in poultry.

Method of Vaccination:

Administer 1 dose (0.5 mL) to each chicken via intramuscular injection.

Withdrawal Period:

6 weeks

Storage Conditions:

Store in a dark place at 2-8°C. Do not freeze.

Packaging:

100Doses/50mL/Bottle, 200Doses/100mL/Bottle, 500Doses/250mL/Bottle, and 1000Doses/500mL/Bottle

Shelf Life:

2 years

Precautions:

- Follow the instructions from and use under supervision by a veterinarian.
- Recover vaccine back to room temperature before use.
- Shake well to mix thoroughly before use.
- Protect from sunlight and high temperatures.
- Vaccinate healthy chickens only.