

BIOFORS ND-IC GEL LASOTA 0.5

Generic Name:

BIOFORS ND-IC Inactivated Vaccine (ND LaSota · IC A&C)

General Description:

BIOFORS ND-IC GEL LASOTA 0.5 contains inactivated Newcastle Disease Virus (NDV) LaSota strain and *Avibacterium paragallinarum* serotype A (IC-A) and *Avibacterium paragallinarum* serotype C (IC-C) mixed with aluminum hydroxide adjuvant. NDV LaSota strain was amplified in embryonated chicken eggs, whereas IC-A and IC-C were amplified in culture media. Allantoic fluids of embryonated chicken eggs containing amplified NDV LaSota strain and culture media containing amplified IC-A and IC-C were harvested and inactivated. Aluminum hydroxide adjuvant was then added. The vaccine has a physical appearance as yellowish-white suspension.

Composition per Dose (0.5mL):

Newcastle Disease Virus LaSota strain	$$ $\geq 10^{8.5}$ EID ₅₀
Avibacterium paragallinarum serotype A	≥10 ^{8.7} CFU
Avibacterium paragallinarum serotype C	$> 10^{8.7}$ CFU

Indication for Use:

Prevention of Newcastle Disease and Infectious Coryza in poultry.

Method of Vaccination:

Administer 1 dose (0.5 mL) to each chicken via intramuscular injection.

Withdrawal Period:

6 weeks

Storage Conditions:

Store in a dark place at 2-8 °C. Do not freeze.

Packaging:

1000Doses/500mL/Bottle

Shelf Life:

2 years

Precautions:

- •Follow the instructions from and use under supervision by a veterinarian. For veterinary use only.
- Protect from light and high temperatures.
- •Allow the vaccine to recover to room temperature before use.
- •Shake well to mix thoroughly before use.
- Vaccinate healthy chickens only.
- Avoid accidental injection into the hands. Seek immediate medical attention if this occurs.

