

## Generic Name:

**BIOFORS ND-MG Inactivated Vaccine** 

### General Description:

BIOFORS ND-MG 0.25 contains inactivated Newcastle Disease Virus (NDV) LaSota and inactivated *Mycoplasma gallisepticum* (MG) in stable oil emulsion. NDV LaSota strain was amplified in embryonated chicken eggs and MG YL-1 strain was amplified in culture medium. Allantoic fluids of embryonated chicken eggs containing amplified NDV LaSota strain and culture medium containing amplified MG YL-1 strain were harvested and oil adjuvant was added. This vaccine has a physical appearance as white oil emulsion.

#### Composition per Dose (0.25mL):

Newcastle Disease Virus LaSota strain $\geq 10$	$^{8.5}$ EID <sub>50</sub>
Mycoplasma gallisepticum YL-1 strain $\geq 1$	0 <sup>7.7</sup> CFU

#### Indication for Use:

Prevention of Newcastle Disease and Mycoplasma gallisepticum infection in poultry.

#### Method of Vaccination:

- 1<sup>st</sup> vaccination: administer 1 dose (0.25 mL) to each chicken via subcutaneous or intramuscular injection.
- 2<sup>nd</sup> vaccination: administer 1 dose (0.25 mL) at 3-2 weeks post-1st vaccination via subcutaneous or intramuscular injection.

#### Withdrawal Period:

8 weeks

<u>Storage Conditions</u>: Store in a dark place at 2-8°C. Do not freeze.

#### Packaging:

200Doses/50mL/Bottle, 400Doses/100mL/Bottle, 1000Doses/250mL/Bottle, and 2000Doses/500mL/Bottle

#### Shelf Life:

2 years

# Precautions:

- •Follow the instructions from and use under supervision by a veterinarian.
- •Recover vaccine to room temperature before use.
- •Shake well to mix thoroughly before use.
- •Protect from sunlight and high temperatures
- •Vaccinate healthy chickens only.
- •Avoid accidental injection into the hands. Seek immediate medical attention if this occurs.

