



BIOFORS ND-IC-MG 0.25

Generic Name:

BIOFORS ND-IC-MG Inactivated Vaccine

General Description:

BIOFORS ND-IC-MG 0.25 contains inactivated Newcastle Disease virus Ishii strain, inactivated *Avibacterium paragallinarum* serotype A and inactivated *Avibacterium paragallinarum* serotype C and inactivated *Mycoplasma gallisepticum* in stable oil emulsion. The product may prevent Newcastle Disease, Infectious Coryza and *Mycoplasma gallisepticum* infections in chicken.

Composition per Dose (0.25mL):

Newcastle Disease Virus Ishii strain.....	$\geq 10^{7.7}$ EID ₅₀
<i>Avibacterium paragallinarum</i> serotype A.....	$\geq 10^{7.6}$ CFU
<i>Avibacterium paragallinarum</i> serotype C.....	$\geq 10^{7.6}$ CFU
<i>Mycoplasma gallisepticum</i> YL-1 strain.....	$\geq 10^{8.0}$ CFU

Indication for Use:

Prevention of Newcastle Disease, Infectious Coryza and *Mycoplasma gallisepticum* infection in chicken.

Method of Vaccination:

Administer 1 dose (0.25mL) to each chicken at 35-70 days old via subcutaneous or intramuscular injection and administer 1 dose (0.25 mL) to each chicken at 90-115 days old via subcutaneous or intramuscular injection.

Withdrawal Period:

8 weeks

Storage Conditions:

Store in a dark place at 2-8°C. Do not freeze.

Packaging:

200Doses/50mL/Bottle, 400Doses/100mL/Bottle, 1000Doses/250mL/Bottle, and 2000Doses/500mL/Bottle

Shelf Life:

2 years

Precautions:

- Follow the instructions from and use under supervision by a veterinarian. For veterinary use only.
- Recover vaccine to room temperature before use.
- Shake well to mix thoroughly before use.
- Protect from sunlight and high temperatures.
- Vaccinate healthy chickens only.
- Avoid accidental injection into the hands. Seek immediate medical attention if this occurs.

