

BIOFORS ND-IC-FC 0.25

Generic Name:

BIOFORS ND-IC-FC Inactivated Vaccine

General Description:

BIOFORS ND-IC-FC 0.25 contains inactivated Newcastle Disease Virus (NDV) Ishii strain, inactivated *Avibacterium paragallinarum* serotype A (IC-A) and inactivated *Avibacterium paragallinarum* serotype C (IC-C), and *Pasteurella multocida* serotype A: 1 strain in a stable oil emulsion. NDV Ishii strain was amplified in embryonated chicken eggs, whereas IC-A, IC-C and *Pasteurella multocida* were amplified in culture media. Allantoic fluids containing amplified NDV Ishii, and culture media containing amplified IC-A, IC-C and *Pasteurella multocida* serotype A: 1 were harvested and inactivated. Oil adjuvant was then added. The vaccine has a physical appearance as white oil emulsion.

Composition per Dose (0.25mL):

Newcastle Disease Virus Ishii strain.....	$\geq 10^{8.5} \text{EID}_{50}$
<i>Avibacterium paragallinarum</i> serotype A.....	$\geq 3 \times 10^7 \text{CFU}$
<i>Avibacterium paragallinarum</i> serotype C.....	$\geq 3 \times 10^7 \text{CFU}$
<i>Pasteurella multocida</i> serotype A: 1.....	$\geq 4 \times 10^8 \text{CFU}$

Indication for Use:

Prevention of infection of Newcastle Disease, Infectious Coryza and Fowl Cholera in poultry.

Method of Vaccination:

This vaccine is given one dose (0.25 mL) per poultry being older than 5 weeks via intramuscular injection.

Withdrawal Period:

8 weeks

Storage Conditions:

Store in a dark place at 2-8°C. Do not freeze.

Packaging:

200Doses/50mL/Bottle, 400Doses/100mL/Bottle, 1000Doses/250mL/Bottle, and 2000Doses/500mL/Bottle

Shelf Life:

2 years

Precautions:

- Follow the instructions from and use under supervision by a veterinarian.
- Recover vaccine to room temperature before use.
- Shake well to mix thoroughly before use.
- Protect from sunlight and high temperatures
- Vaccinate healthy chickens only.
- Avoid accidental injection into the hands of vaccinators and seek immediate medical attention if this occurs.